



Patient's Name: \_\_\_\_\_ DOB: \_\_\_\_\_ DATE: \_\_\_\_\_

**YALE-BROWN OBSESSIVE COMPULSIVE SCALE (Y-BOCS)\*****Questions 1 to 5 are about your obsessive thoughts**

Obsessions are unwanted ideas, images or impulses that intrude on thinking against your wishes and efforts to resist them. They usually involve themes of harm, risk and danger. Common obsessions are excessive fears of contamination; recurring doubts about danger, extreme concern with order, symmetry, or exactness; fear of losing important things.

Please answer each question by circling the appropriate number.

**1. TIME OCCUPIED BY OBSESSIVE THOUGHTS** SCORE \_\_\_\_\_

How much of your time is occupied by obsessive thoughts?

- |   |   |  |
|---|---|--|
| 0 | = | None   |
| 1 | = | Less than 1 hr/day or occasional occurrence                    |
| 2 | = | 1 to 3 hrs/day or frequent                                     |
| 3 | = | Greater than 3 and up to 8 hrs/day or very frequent occurrence |
| 4 | = | Greater than 8 hrs/day or nearly constant occurrence           |

**2. INTERFERENCE DUE TO OBSESSIVE THOUGHTS** SCORE \_\_\_\_\_

How much do your obsessive thoughts interfere with your work, school, social, or other important role functioning? Is there anything that you don't do because of them?

- |   |   |   |
|---|---|---|
| 0 | = | None  |
| 1 | = | Slight interference with social or other activities, but overall performance not impaired |
| 2 | = | Definite interference with social or occupational performance, but still manageable       |
| 3 | = | Causes substantial impairment in social or occupational performance                       |
| 4 | = | Incapacitating  |

**3. DISTRESS ASSOCIATED WITH OBSESSIVE THOUGHTS** SCORE \_\_\_\_\_

How much distress do your obsessive thoughts cause you?

- |   |   |                                      |
|---|---|--------------------------------------|
| 0 | = | None                                 |
| 1 | = | Not too disturbing                   |
| 2 | = | Disturbing, but still manageable     |
| 3 | = | Very disturbing                      |
| 4 | = | Near constant and disabling distress |

**4. RESISTANCE AGAINST OBSESSIONS** SCORE \_\_\_\_\_

How much of an effort do you make to resist the obsessive thoughts? How often do you try to disregard or turn your attention away from these thoughts as they enter your mind?

- |   |   |  |
|---|---|--|
| 0 | = | Try to resist all the time   |
| 1 | = | Try to resist most of the time   |
| 2 | = | Make some effort to resist   |
| 3 | = | Yield to all obsessions without attempting to control them, but with some reluctance |
| 4 | = | Completely and willingly yield to all obsessions                                     |

## 5. DEGREE OF CONTROL OVER OBSESSIVE THOUGHTS

SCORE \_\_\_\_\_

How much control do you have over your obsessive thoughts? How successful are you in stopping or diverting your obsessive thinking? Can you dismiss them?

- |   |   |   |
|---|---|---|
| 0 | = | Complete control  |
| 1 | = | Usually able to stop or divert obsessions with some effort and concentration                      |
| 2 | = | Sometimes able to stop or divert obsessions   |
| 3 | = | Rarely successful in stopping or dismissing obsessions, can only divert attention with difficulty |
| 4 | = | Obsessions are completely involuntary, rarely able to even momentarily alter obsessive thinking.  |

The next several questions are about your compulsive behaviors.

Compulsions are urges that people have to do something to lessen feelings of anxiety or other discomfort. Often they do repetitive, purposeful, intentional behaviors called rituals. The behavior itself may seem appropriate but it becomes a ritual when done to excess. Washing, checking, repeating, straightening, hoarding and many other behaviors can be rituals. Some rituals are mental. For example, thinking or saying things over and over under your breath.

## 6. TIME SPENT PERFORMING COMPULSIVE BEHAVIORS

SCORE \_\_\_\_\_

How much time do you spend performing compulsive behaviors? How much longer than most people does it take to complete routine activities because of your rituals? How frequently do you do rituals?

- |   |   |   |
|---|---|---|
| 0 | = | None  |
| 1 | = | Less than 1 hr/day or occasional performance of compulsive behaviors                              |
| 2 | = | From 1 to 3 hrs/day, or frequent performance of compulsive behaviors                              |
| 3 | = | More than 3 and up to 8 hrs/day, or very frequent performance of compulsive behaviors             |
| 4 | = | More than 8 hrs/day, or near constant performance of compulsive behaviors (too numerous to count) |

## 7. INTERFERENCE DUE TO COMPULSIVE BEHAVIORS

SCORE \_\_\_\_\_

How much do your compulsive behaviors interfere with your work, school, social, or other important role functioning? Is there anything that you don't do because of the compulsions?

- |   |   |   |
|---|---|---|
| 0 | = | None  |
| 1 | = | Slight interference with social or other activities, but overall performance not impaired |
| 2 | = | Definite interference with social or occupational performance, but still manageable       |
| 3 | = | Causes substantial impairment in social or occupational performance                       |
| 4 | = | Incapacitating  |

## 8. DISTRESS ASSOCIATED WITH COMPULSIVE BEHAVIOR

SCORE \_\_\_\_\_

How would you feel if prevented from performing your compulsion(s)? How anxious would you become?

- |   |   |  |
|---|---|--|
| 0 | = | None   |
| 1 | = | Only slightly anxious if compulsions prevented                               |
| 2 | = | Anxiety would mount but remain manageable if compulsions prevented           |
| 3 | = | Prominent and very disturbing increase in anxiety if compulsions interrupted |
| 4 | = | Incapacitating anxiety from any intervention aimed at modifying activity     |

## 9. RESISTANCE AGAINST COMPULSIONS

SCORE \_\_\_\_\_

How much of an effort do you make to resist the compulsions?

- |   |   |  |
|---|---|--|
| 0 | = | Always try to resist   |
| 1 | = | Try to resist most of the time   |
| 2 | = | Make some effort to resist   |
| 3 | = | Yield to almost all compulsions without attempting to control them, but with some reluctance |
| 4 | = | Completely and willingly yield to all compulsions  |

## 10. DEGREE OF CONTROL OVER COMPULSIVE BEHAVIOR

SCORE \_\_\_\_\_

How strong is the drive to perform the compulsive behavior? How much control do you have over the compulsions?

- |   |   |   |
|---|---|---|
| 0 | = | Complete control  |
| 1 | = | Pressure to perform the behavior but usually able to exercise voluntary control over it   |
| 2 | = | Strong pressure to perform behavior, can control it only with difficulty  |
| 3 | = | Very strong drive to perform behavior, must be carried to completion, can only delay with difficulty                              |
| 4 | = | Drive to perform behavior experienced as completely involuntary and overpowering, rarely able to even momentarily delay activity. |

---

 TOTAL SCORE \_\_\_\_\_

# Y-BOCS Symptom Checklist

Instructions: Generate a *Target Symptoms List* from the attached Y-BOCS Symptom Checklist by asking the patient about specific obsessions and compulsions. Check all that apply. Distinguish between current and past symptoms. Mark principal symptoms with a "p". These will form the basis of the *Target Symptoms List*. Items marked may "\*" or may not be an OCD phenomena.

Current Past

## AGGRESSIVE OBSESSIONS

- |     |     |  |
|-----|-----|--|
| ___ | ___ | Fear might harm self   |
| ___ | ___ | Fear might harm others   |
| ___ | ___ | Violent or horrific images   |
| ___ | ___ | Fear of blurting out obscenities or insults  |
| ___ | ___ | Fear of doing something else embarrassing*   |
| ___ | ___ | Fear will act on unwanted impulses (e.g., to stab friend)                              |
| ___ | ___ | Fear will steal things   |
| ___ | ___ | Fear will harm others because not careful enough (e.g. hit/run motor vehicle accident) |
| ___ | ___ | Fear will be responsible for something else terrible happening (e.g., fire, burglary)  |

Other: \_\_\_\_\_

## CONTAMINATION OBSESSIONS

- |     |     |  |
|-----|-----|--|
| ___ | ___ | Concerns or disgust w/ with bodily waste or secretions (e.g., urine, feces, saliva)      |
| ___ | ___ | Concern with dirt or germs   |
| ___ | ___ | Excessive concern with environmental contaminants (e.g. asbestos, radiation toxic waste) |
| ___ | ___ | Excessive concern with household items (e.g., cleansers solvents)                        |
| ___ | ___ | Excessive concern with animals (e.g., insects)   |
| ___ | ___ | Bothered by sticky substances or residues  |
| ___ | ___ | Concerned will get ill because of contaminant  |
| ___ | ___ | Concerned will get others ill by spreading contaminant (Aggressive)                      |
| ___ | ___ | No concern with consequences of contamination other than how it might feel               |

## SEXUAL OBSESSIONS

- |     |     |  |
|-----|-----|--|
| ___ | ___ | Forbidden or perverse sexual thoughts. images. or impulses |
| ___ | ___ | Content involves children or incest                        |
| ___ | ___ | Content involves homosexuality*                            |
| ___ | ___ | Sexual behavior towards others (Aggressive)*               |
| ___ | ___ | Other: _____   |

## HOARDING/SAVING OBSESSIONS

(distinguish from hobbies and concern with objects of monetary or sentimental value)

## RELIGIOUS OBSESSIONS (Scrupulosity)

- |     |     |   |
|-----|-----|---|
| ___ | ___ | Concerned with sacrilege and blasphemy    |
| ___ | ___ | Excess concern with right/wrong, morality |
| ___ | ___ | Other: _____                              |

## OBSESSION WITH NEED FOR SYMMETRY OR EXACTNESS

- |     |     |  |
|-----|-----|--|
| ___ | ___ | Accompanied by magical thinking (e.g., concerned that another will have accident dent unless less things are in the right place) |
| ___ | ___ | Not accompanied by magical thinking  |

## MISCELLANEOUS OBSESSIONS

- |     |     |  |
|-----|-----|--|
| ___ | ___ | Need to know or remember                   |
| ___ | ___ | Fear of saying certain things              |
| ___ | ___ | Fear of not saying just the right thing    |
| ___ | ___ | Fear of losing things                      |
| ___ | ___ | Intrusive (nonviolent) images              |
| ___ | ___ | Intrusive nonsense sounds, words, or music |
| ___ | ___ | Bothered by certain sounds/noises*         |
| ___ | ___ | Lucky/unlucky numbers                      |
| ___ | ___ | Colors with special significance           |
| ___ | ___ | 3 superstitious fears                      |
| ___ | ___ | Other: _____                               |

Current Past

## SOMATIC OBSESSIONS

- |     |     |  |
|-----|-----|--|
| ___ | ___ | Concern with illness or disease*   |
| ___ | ___ | Excessive concern with body part or aspect of Appearance (eg., dysmorphophobia)* |
| ___ | ___ | Other _____  |

## CLEANING/WASHING COMPULSIONS

- |     |     |   |
|-----|-----|---|
| ___ | ___ | Excessive or ritualized handwashing   |
| ___ | ___ | Excessive or ritualized showering, bathing, toothbrushing grooming, or toilet routine Involves cleaning of household items or other inanimate objects |
| ___ | ___ | Other measures to prevent or remove contact with contaminants   |
| ___ | ___ | Other _____   |

## CHECKING COMPULSIONS

- |     |     |  |
|-----|-----|--|
| ___ | ___ | Checking locks, stove, appliances etc.         |
| ___ | ___ | Checking that did rot/will not harm others     |
| ___ | ___ | Checking that did not/will not harm self       |
| ___ | ___ | Checking that nothing terrible did/will happen |
| ___ | ___ | Checking that did not make mistake             |
| ___ | ___ | Checking tied to somatic obsessions            |
| ___ | ___ | Other: _____                                   |

## REPEATING RITUALS

- |     |     |   |
|-----|-----|---|
| ___ | ___ | Rereading or rewriting  |
| ___ | ___ | Need to repeat routine activities jog, in/out door, up/down from chair) |
| ___ | ___ | Other _____   |

## COUNTING COMPULSIONS

\_\_\_\_\_

## ORDERING/ARRANGING COMPULSIONS

\_\_\_\_\_

## HOARDING/COLLECTING COMPULSIONS

(distinguish from hobbies and concern with objects of monetary or sentimental value (e.g., carefully reads junk mail, piles up old newspapers, sorts through garbage, collects useless objects.)

\_\_\_\_\_

## MISCELLANEOUS COMPULSIONS

- |     |     |   |
|-----|-----|---|
| ___ | ___ | Mental rituals (other than checking/counting)   |
| ___ | ___ | Excessive listmaking  |
| ___ | ___ | Need to tell, ask, or confess   |
| ___ | ___ | Need to touch, tap, or rub*   |
| ___ | ___ | Rituals involving blinking or staring*  |
| ___ | ___ | Measures (not checking) to prevent: harm to self - harm to others terrible consequences |
| ___ | ___ | Ritualized eating behaviors*  |
| ___ | ___ | Superstitious behaviors   |
| ___ | ___ | Trichotillomania *  |
| ___ | ___ | Other self-damaging or self-mutilating behaviors*                                       |
| ___ | ___ | Other _____   |

Adapted from Goodman, W.K., Price, L.H., Rasmussen, S.A. et al.:  
"The Yale-Brown Obsessive Compulsive Scale."  
Arch Gen Psychiatry 46:1006-1011,1989

**Original:** 10/1/86  
**First Revision:** 3/1/90  
**Second Revision:** 5/1/91  
**Third Revision:** 5/1/93  
**Fourth Revision:** 6/17/99  
**Fifth Revision:** 10/04/07

**CHILDREN'S  
YALE-BROWN  
OBSESSIVE COMPULSIVE SCALE  
(CY-BOCS)**

**DEVELOPED BY**

**WAYNE K. GOODMAN, M.D.<sup>1</sup>  
LAWRENCE SCAHILL, MSN, PhD<sup>2</sup>  
LAWRENCE H. PRICE, M.D.<sup>3</sup>  
STEVEN A. RASMUSSEN, M.D.<sup>3</sup>  
MARK A. RIDDLE, M.D.<sup>4</sup>  
JUDITH L. RAPOPORT, M.D.<sup>5</sup>**

**NATIONAL INSTITUTE OF MENTAL HEALTH<sup>1</sup>**

**THE CHILD STUDY CENTER<sup>2</sup>  
YALE UNIVERSITY SCHOOL OF MEDICINE**

**DEPARTMENT OF PSYCHIATRY<sup>3</sup>  
BROWN UNIVERSITY SCHOOL OF MEDICINE**

**CHILD PSYCHIATRY DIVISION<sup>4</sup>  
JOHNS HOPKINS SCHOOL OF MEDICINE  
and**

**CHILD PSYCHIATRY BRANCH<sup>5</sup>  
NATIONAL INSTITUTE OF MENTAL HEALTH**

**Investigators interested in using this rating scale should contact Lawrence Scahill, M.S.N., Ph.D., at the Yale Child Study Center, P.O. Box 207900, New Haven, CT 06520 or Wayne Goodman, M.D., at the National Institute of Mental Health, Bethesda, MD.**

Scahill, L., Riddle, M.A., McSwiggin-Hardin, M., Ort, S.I., King, R.A., Goodman, W.K., Cicchetti, D. & Leckman, J.F. (1997). Children's Yale-Brown Obsessive Compulsive Scale: reliability and validity. *J Am Acad Child Adolesc Psychiatry*, 36(6):844-852.

## **GENERAL INSTRUCTIONS**

### **Overview:**

This scale is designed to rate the severity of obsessive and compulsive symptoms in children and adolescents, ages 6 to 17 years. It can be administered by a clinician or trained interviewer in a semi-structured fashion. In general, the ratings depend on the child's and parent's report; however, the final rating is based on the clinical judgement of the interviewer. Rate the characteristics of each item over the prior week up until, and including, the time of the interview. Scores should reflect the average of each item for the entire week, unless otherwise specified.

### **Informants:**

Information should be obtained by interviewing the parent(s) (or guardian) and the child together. Sometimes, however, it may also be useful to interview the child or parent alone. Interviewing strategy may vary depending on the age and developmental level of the child or adolescent. All information should be combined to estimate the score for each item. Whenever the CY-BOCS is administered more than once to the same child, as in a medication trial, consistent reporting can be ensured by having the same informant(s) present at each rating session.

### **Definitions:**

Before proceeding with the questions, define "obsessions" and "compulsions" for the child and primary caretaker as follows (sometimes, particularly with younger children, the interviewer may prefer using the terms "worries" and "habits"):

"OBSESSIONS: are thoughts, ideas, or pictures that keep coming into your mind even though you do not want them to. They may be unpleasant, silly or embarrassing."

"AN EXAMPLE OF AN OBSESSION IS: the repeated thought that germs or dirt are harming you or other people, or that something unpleasant might happen to you or someone in your family or someone special to you. These are thoughts that keep coming back, over and over again."

"COMPULSIONS: are things that you feel you have to do although you may know that they do not make sense. Sometimes you may try to stop from doing them but this might not be possible. You might feel worried or angry or frustrated until you have finished what you have to do."

"AN EXAMPLE OF A COMPULSION IS: the need to wash your hands over and over again even though they are not really dirty, or the need to count up to a certain number while you do certain things."

"Do you have any questions about what these words called obsessions and compulsions mean?"

### **Symptom Specificity and Continuity:**

In some cases, it may be difficult to delineate obsessions and compulsions from other closely related symptoms such as phobias, anxious worries, depressive ruminations or complex tics. Separate assessment of these symptoms may be necessary. Although potentially difficult, the delineation of obsessions and compulsions from these closely related symptoms is an essential task of the interviewer. (A full discussion of how to make this determination is beyond the scope and purpose of this introduction.) Items marked with an asterisk are items where this delineation may be especially troublesome.

Once the interviewer has decided whether or not a particular symptom will be included as an obsession or compulsion on the checklist, every effort should be made to maintain consistency in subsequent rating(s). In a treatment study with multiple ratings over time, it may be useful to review the initial Target Symptom Checklist (see below) at the beginning of subsequent ratings (prior severity scores should not be reviewed).

### **Procedure:**

Symptom Checklist: After reviewing with the child and parent(s) the definitions of obsessions and compulsions, the interview should proceed with a detailed inquiry about the child's symptoms using the Compulsions Checklist and Obsessions Checklist as guides. It may not be necessary to ask about each and every item on the checklist, but each symptom area should be covered to ensure that symptoms are not missed. For most children and adolescents, it is usually easier to begin with compulsions (pages 9 and 10).

Target Symptom List: After the Compulsions Checklist is complete, list the four most severe compulsions on the Target Symptom List on page 10. Repeat this process, listing the most severe obsessions, on the Target Symptom List on page 5.

Severity Rating: After completing the Checklist and Target Symptom List for compulsions, inquire about the severity items: Time Spent, Distress, Resistance, Interference, and Degree of Control (questions 6 through 10 on pages 11 through 13). There are examples of probe questions for each item. Ratings for these items should reflect interviewer's best estimate from all available information from the past week, with special emphasis on the Target Symptoms. Repeat the above procedure for obsessions (Pages 4 through 8). Finally, inquire about and rate questions 11 through 19 on pages 14 and 18. Scores can be recorded on the scoring sheet on page 19. All ratings should be in whole integers.

### **Scoring:**

All 19 items are rated, but only items 1-10 are used to determine the total score. The **total CY-BOCS score** is the sum of items 1-10; the **obsession and compulsion subtotals** are the sums of items 1-5 and 6-10, respectively. At this time, items 1A and 6A are not being used in the scoring.

Items 17 (global severity) and 18 (global improvement) are adapted from the Clinical Global Impression Scale (Guy, W., 1976) to provide measures of overall functional impairment associated with the presence of obsessive-compulsive symptoms.

Name \_\_\_\_\_ Date \_\_\_\_\_

### CY-BOCS OBSESSIONS CHECKLIST

Check all items that apply (Item marked "\*" may or not be OCD phenomena.)

**Current**   **Past**   **Contamination Obsessions**

- |       |       |  |
|-------|-------|--|
| _____ | _____ | Concern with dirt, germs, certain illnesses (e.g., AIDS)                                   |
| _____ | _____ | Concerns or disgust with bodily waste or secretions (e.g., urine, feces, saliva)           |
| _____ | _____ | Excessive concern with environmental contaminants (e.g., asbestos, radiation, toxic waste) |
| _____ | _____ | Excessive concern with household items (e.g., cleaners, solvents)                          |
| _____ | _____ | Excessive concern about animals/insects  |
| _____ | _____ | Excessively bothered by sticky substances or residues                                      |
| _____ | _____ | Concerned will get ill because of contaminant  |
| _____ | _____ | Concerned will get others ill by spreading contaminant (aggressive)                        |
| _____ | _____ | No concern with consequences of contamination other than how it might feel *               |
| _____ | _____ | Other (Describe) _____   |

**Aggressive Obsessions**

- |       |       |   |
|-------|-------|---|
| _____ | _____ | Fear might harm self  |
| _____ | _____ | Fear might harm others  |
| _____ | _____ | Fear harm will come to self   |
| _____ | _____ | Fear harm will come to others (may be because something child did or did not do)            |
| _____ | _____ | Violent or horrific images  |
| _____ | _____ | Fear of blurting out obscenities or insults   |
| _____ | _____ | Fear of doing something else embarrassing *   |
| _____ | _____ | Fear will act on unwanted impulses (e.g. to stab a family member)                           |
| _____ | _____ | Fear will steal things  |
| _____ | _____ | Fear will be responsible for something else terrible happening (e.g. fire, burglary, flood) |
| _____ | _____ | Other (Describe) _____  |

**Sexual Obsessions**

[Are you having any sexual thoughts? If yes, are they routine or are they repetitive thoughts that you would rather not have or find disturbing? If yes, are they:]

- |       |       |   |
|-------|-------|---|
| _____ | _____ | Forbidden or perverse sexual thoughts, images, impulses |
| _____ | _____ | Content involves homosexuality *                        |
| _____ | _____ | Sexual behavior towards others (Aggressive)             |
| _____ | _____ | Other (Describe) _____                                  |

**Hoarding/Saving Obsessions**

- |       |       |                        |
|-------|-------|------------------------|
| _____ | _____ | Fear of losing things  |
| _____ | _____ | Other (Describe) _____ |

**Magical Thoughts/Superstitious Obsessions**

- |       |       |                                      |
|-------|-------|--------------------------------------|
| _____ | _____ | Lucky/unlucky numbers, colors, words |
| _____ | _____ | Other (Describe) _____               |



**Current   Past      Somatic Obsessions**

\_\_\_\_\_    \_\_\_\_\_    Excessive concern with illness or disease \*

\_\_\_\_\_    \_\_\_\_\_    Excessive concern with body part or aspect of appearance (e.g., dysmorphophobia) \*

\_\_\_\_\_    \_\_\_\_\_    Other (Describe) \_\_\_\_\_

**Religious Obsessions (Scrupulosity)**

\_\_\_\_\_    \_\_\_\_\_    Excessive concern or fear of offending religious objects (God)

\_\_\_\_\_    \_\_\_\_\_    Excessive concern with right/wrong, morality

\_\_\_\_\_    \_\_\_\_\_    Other (Describe) \_\_\_\_\_

**Miscellaneous Obsessions**

\_\_\_\_\_    \_\_\_\_\_    The need to know or remember

\_\_\_\_\_    \_\_\_\_\_    Fear of saying certain things

\_\_\_\_\_    \_\_\_\_\_    Fear of not saying just the right thing

\_\_\_\_\_    \_\_\_\_\_    Intrusive (non-violent) images

\_\_\_\_\_    \_\_\_\_\_    Intrusive sounds, words, music, or numbers

\_\_\_\_\_    \_\_\_\_\_    Other (Describe) \_\_\_\_\_

**TARGET SYMPTOM LIST FOR OBSESSIONS**

**Obsessions (Describe, listing by order of severity, with #1 being the most severe, #2 the second most severe, etc.):**

1. \_\_\_\_\_
2. \_\_\_\_\_
3. \_\_\_\_\_
4. \_\_\_\_\_

**QUESTIONS ON OBSESSIONS (ITEMS 1-5)** "I AM NOW GOING TO ASK YOU QUESTIONS ABOUT THE THOUGHTS YOU CANNOT STOP THINKING ABOUT." (Review for the informant(s) the Target Symptoms and refer to them while asking questions 1-5).

1. Time Occupied by Obsessive Thoughts

- How much time do you spend thinking about these things?  
(When obsessions occur as brief, intermittent intrusions, it may be impossible to assess time occupied by them in terms of total hours. In such cases, estimate time by determining how frequently they occur. Consider both the number of times the intrusions occur and how many hours of the day are affected).
- How frequently do these thoughts occur?  
[Exclude ruminations and preoccupations which, unlike obsessions, are ego-syntonic and rational (but exaggerated).]

0 - NONE

1 - MILD                      less than 1 hr/day or occasional intrusion

2 - MODERATE            1 to 3 hrs/day or frequent intrusion

3 - SEVERE                greater than 3 and up to 8 hrs/day or very frequent intrusion

4 - EXTREME              greater than 8 hrs/day or near constant intrusion

1B. Obsession-free Interval (not included in total score)

- On average, what is the longest amount of time per day that you are not bothered by obsessive thoughts?

0 - NONE

1 - MILD                      long symptom free intervals, more than 8 consecutive hrs/day symptom-free

2 - MODERATE            moderately long symptom-free intervals, more than 3 and up to 8 hrs/day

3 - SEVERE                brief symptom-free intervals, from 1 to 3 consecutive hrs/day symptom-free

4 - EXTREME              less than 1 consecutive hr/day symptom free

2. Interference due to Obsessive Thoughts

- How much do these thoughts get in the way of school or doing things with friends?
- Is there anything that you don't do because of them?  
(If currently not in school determine how much performance would be affected if patient were in school.)

0 - NONE

1 - MILD                      slight interference with social or school activities, overall performance not impaired

2 - MODERATE            definite interference with social or school performance, but still manageable

3 - SEVERE                causes substantial impairment in social or school performance

4 - EXTREME              incapacitating

### 3. Distress Associated with Obsessive Thoughts

- How much do these thoughts bother or upset you?

(Only rate anxiety/frustration that seems triggered by obsessions, not generalized anxiety or anxiety associated with other symptoms.)

0 - NONE

1 - MILD                      infrequent, and not too disturbing

2 - MODERATE              frequent, and disturbing, but still manageable

3 - SEVERE                  very frequent, and very disturbing

4 - EXTREME                near constant, and disabling distress/frustration

### 4. Resistance Against Obsessions

- How hard do you try to stop the thoughts or ignore them?

(Only rate effort made to resist, not success or failure in actually controlling the obsessions. How much patient resists the obsessions may or may not correlate with their ability to control them. Note that this item does not directly measure the severity of the intrusive thoughts; rather it rates a manifestation of health, i.e., the effort the patient makes to counteract the obsessions. Thus, the more the patient tries to resist, the less impaired is this aspect of his functioning. If the obsessions are minimal, the patient may not feel the need to resist them. In such cases, a rating of "0" should be given.)

0 – NONE                      makes an effort to always resist, or symptoms so minimal doesn't need to actively resist.

1 - MILD                      tries to resist most of the time

2 - MODERATE              makes some effort to resist

3 - SEVERE                  yields to all obsessions without attempting to control them, but does so with some reluctance

4 – EXTREME                completely and willingly yields to all obsessions

### 5. Degree of Control Over Obsessive Thoughts

- When you try to fight the thoughts, can you beat them?

- How much control do you have over the thoughts?

(In contrast to the preceding item on resistance, the ability of the patient to control his obsessions is more closely related to the severity of the intrusive thoughts.)

0 - COMPLETE CONTROL

1 - MUCH CONTROL              usually able to stop or divert obsessions with some effort and concentration.

2 - MODERATE CONTROL      sometimes able to stop or divert obsessions

3 - LITTLE CONTROL            rarely successful in stopping obsessions, can only divert attention with difficulty

4 - NO CONTROL                experienced as completely involuntary, rarely able to even momentarily divert thinking

Name \_\_\_\_\_ Date \_\_\_\_\_

## CY-BOCS COMPULSIONS CHECKLIST

Check all items that apply (Item marked "\*" may or not be OCD phenomena.)

---

### **Current Past Washing/Cleaning Compulsions**

- |       |       |  |
|-------|-------|--|
| _____ | _____ | Excessive or ritualized handwashing  |
| _____ | _____ | Excessive or ritualized showering, bathing, toothbrushing, grooming, or toilet routine |
| _____ | _____ | Excessive cleaning of items; such as personal clothes or important objects             |
| _____ | _____ | Other measures to prevent or remove contact with contaminants                          |
| _____ | _____ | Other (Describe) _____   |

### **Checking Compulsions**

- |       |       |   |
|-------|-------|---|
| _____ | _____ | Checking locks, toys, school books/items, etc.                  |
| _____ | _____ | Checking associated with getting washed, dressed, or undressed. |
| _____ | _____ | Checking that did not/will not harm others                      |
| _____ | _____ | Checking that did not/will not harm self                        |
| _____ | _____ | Checking that nothing terrible did/will happen                  |
| _____ | _____ | Checking that did not make mistake                              |
| _____ | _____ | Checking tied to somatic obsessions                             |
| _____ | _____ | Other (Describe) _____  |

### **Repeating Rituals**

- |       |       |   |
|-------|-------|---|
| _____ | _____ | Rereading, erasing, or rewriting  |
| _____ | _____ | Need to repeat routine activities (e.g. in/out doors, up/down from chair) |
| _____ | _____ | Other (Describe) _____  |

### **Counting Compulsions**

- |       |       |                                       |
|-------|-------|---------------------------------------|
| _____ | _____ | Objects, certain numbers, words, etc. |
| _____ | _____ | Describe: _____                       |

### **Ordering/Arranging**

- |       |       |   |
|-------|-------|---|
| _____ | _____ | Need for symmetry/evening up (e.g., lining items up a certain way or arranging personal items in specific patterns) |
| _____ | _____ | Other (Describe) _____  |

### **Hoarding/Saving Compulsion**

- |       |       |  |
|-------|-------|--|
| _____ | _____ | [distinguish from hobbies and concern with objects of monetary or sentimental value] |
| _____ | _____ | Difficulty throwing things away, saving bits of paper, string, etc.                  |
| _____ | _____ | Other (Describe) _____   |

### **Excessive Games/Superstitious Behaviors**

- |       |       |   |
|-------|-------|---|
| _____ | _____ | [distinguish from age appropriate magical games]  |
| _____ | _____ | (e.g., array of behavior, such as stepping over certain spots on a floor, touching an object/self certain number of times as a routine game to avoid something bad from happening.) |
| _____ | _____ | Other (Describe) _____  |

**Current   Past   Rituals Involving Other Persons**

The need to involve another person (usually a parent) in ritual (e.g., asking a parent to repeatedly answer the same question, making mother perform certain meal time-rituals involving specific utensils).\*

\_\_\_\_\_   Other (Describe) \_\_\_\_\_

**Miscellaneous Compulsions**

\_\_\_\_\_   Mental rituals (other than checking/counting)

\_\_\_\_\_   Need to tell, ask, or confess

\_\_\_\_\_   Measures (not checking) to prevent harm to self\_\_\_\_; harm to others\_\_\_\_; terrible consequences \_\_\_\_

\_\_\_\_\_   Ritualized eating behaviors \*

\_\_\_\_\_   Excessive list making \*

\_\_\_\_\_   Need to touch, tap, rub \*

\_\_\_\_\_   Need to do things (e.g., touch or arrange) until it feels just right) \*

\_\_\_\_\_   Rituals involving blinking or staring \*

\_\_\_\_\_   Trichotillomania (hair-pulling) \*

\_\_\_\_\_   Other self-damaging or self-mutilating behaviors \*

\_\_\_\_\_   Other (Describe) \_\_\_\_\_

**TARGET SYMPTOM LIST FOR COMPULSIONS**

**Compulsions (Describe, listing by order of severity, with #1 being the most severe, #2 second most severe, etc.):**

1. \_\_\_\_\_

2. \_\_\_\_\_

3. \_\_\_\_\_

4. \_\_\_\_\_

**QUESTIONS ON COMPULSIONS (ITEMS 6-10)** "I AM NOW GOING TO ASK YOU QUESTIONS ABOUT THE HABITS YOU CAN'T STOP." (Review for the informant(s) the Target Symptoms and refer to them while asking questions 6-10).

**6A. Time Spent Performing Compulsive Behaviors**

- How much time do you spend doing these things?
- How much longer than most people does it take to complete your usual daily activities because of the habits?

(When compulsions occur as brief, intermittent behaviors, it may be impossible to assess time spent performing them in terms of total hours. In such cases, estimate time by determining how frequently they are performed. Consider both the number of times compulsions are performed and how many hours of the day are affected.)

- How often do you do these habits?

[In most cases compulsions are observable behaviors (e.g., handwashing), but there are instances in which compulsions are not observable (e.g., silent checking).]

0 - NONE

1 - MILD (spends less than 1 hr/day performing compulsions), or occasional performance of compulsive behaviors

2 - MODERATE (spends from 1 to 3 hrs/day performing compulsions), or frequent performance of compulsive behaviors

3 - SEVERE (spends more than 3 and up to 8 hrs/day performing compulsions), or very frequent performance of compulsive behaviors

4 - EXTREME (spends more than 8 hrs/day performing compulsions), or near constant performance of compulsive behaviors (too numerous to count).

**6B. Compulsion-free Interval**

- How long can you go without performing compulsive behavior?

[If necessary ask: What is the longest block of time in which (your habits) compulsions are absent?]

0 - NO SYMPTOMS

1 - MILD long symptom-free interval, more than 8 consecutive hrs/day symptom-free

2 - MODERATE moderately long symptom-free interval, more than 3 and up to 8 consecutive hrs/day symptom-free.

3 - SEVERE short symptom-free interval, from 1 to 3 consecutive hrs/day symptom free

4 - EXTREME less than 1 consecutive hr/day symptom-free

### 7. Interference due to Compulsive Behaviors

- How much do these habits get in the way of school or doing things with friends?
- Is there anything you don't do because of them?  
(If currently not in school, determine how much performance would be affected if patient were in school.)

0 - NONE

1 - MILD                    slight, interference with social or school activities, but overall performance not impaired

2 – MODERATE            definite interference with social or school performance, but still manageable

3 - SEVERE                causes substantial impairment in social or school performance

4 - EXTREME              incapacitating

### 8. Distress Associated with Compulsive Behavior

- How would you feel if prevented from carrying out your habits?
- How upset would you become?  
(Rate degree of distress/frustration patient would experience if performance of the compulsion were suddenly interrupted without reassurance offered. In most, but not all cases, performing compulsions reduces anxiety /frustration.)
- How upset do you get while carrying out your habits until you are satisfied?

0 - NONE

1 - MILD                    only slightly anxious/frustrated if compulsions prevented, or only slight anxiety/frustration during performance of compulsions.

2 - MODERATE            reports that anxiety/frustration would mount but remain manageable if compulsions prevented. Anxiety/frustration increases but remains manageable during performance of compulsions.

3 - SEVERE                prominent and very disturbing increase in anxiety/frustration if compulsions interrupted. Prominent and very disturbing increase in anxiety /frustration during performance of compulsions.

4 - EXTREME              incapacitating anxiety/frustration from any intervention aimed at modifying activity. Incapacitating anxiety/frustration develops during performance of compulsions.

## 9. Resistance Against Compulsions

- How much do you try to fight the habits?

(Only rate effort made to resist, not success or failure in actually controlling the compulsions. How much the patient resists the compulsions may or may not correlate with his ability to control them. Note that this item does not directly measure the severity of the compulsions; rather it rates a manifestation of health, i.e., the effort the patient makes to counteract the compulsions. Thus, the more the patient tries to resist, the less impaired is this aspect of their functioning. If the compulsions are minimal, the patient may not feel the need to resist them. In such cases, a rating of "0" should be given.)

0 - NONE	Makes an effort to always resist, or symptoms so minimal doesn't need to actively resist.
1 - MILD	Tries to resist most of the time.
2 - MODERATE	Makes some effort to resist
3 - SEVERE	Yields to almost all compulsions without attempting to control them, but does so with some reluctance.
4 - EXTREME	completely and willingly yields to all compulsions

## 10. Degree of Control over Compulsive Behavior

- How strong is the feeling that you have to carry out the habit(s)?

- When you try to fight them what happens?

(For the advanced child ask:)

- How much control do you have over the habits?

(In contrast to the preceding item on resistance, the ability of the patient to control his compulsions is closely related to the severity of the compulsions.)

0 - COMPLETE CONTROL

1 - MUCH CONTROL	experiences pressure to perform the behavior, but usually able to exercise voluntary control over it
2 - MODERATE CONTROL	moderate control, strong pressure to perform behavior, can control it only with difficulty
3 - LITTLE CONTROL	little control, very strong drive to perform behavior, must be carried to completion, can only delay with difficulty
4 - NO CONTROL	no control, drive to perform behavior experienced as completely involuntary and overpowering, rarely able to delay activity (even momentarily)



# CHILDREN'S YALE-BROWN OBSESSIVE COMPULSIVE SCALE

CYBOCS TOTAL (add items 1-10) \_\_\_\_\_

Patient Name \_\_\_\_\_ Date \_\_\_\_\_ Patient ID \_\_\_\_\_  
 Rater \_\_\_\_\_

1. TIME SPENT ON OBSESIONS	0	1	2	3	4
1b. OBSESSION-FREE INTERVAL	No Symptoms	Long	Moderately Long	Short	Extremely Short
(do not add to subtotal or total score)	0	1	2	3	4
2. INTERFERENCE FROM OBSESSIONS	0	1	2	3	4
3. DISTRESS OF OBSESSIONS	0	1	2	3	4
4. RESISTANCE	Always resists 0	1	2	3	Completely yields 4
5. CONTROL OVER OBSESSIONS	Complete control 0	Much control 1	Moderate control 2	Little control 3	No control 4
OBSESSION SUBTOTAL (add items 1-5)					[      ]
6. TIME SPENT ON COMPULSIONS	0	1	2	3	4
6b. COMPULSION-FREE INTERVAL	No Symptoms	Long	Moderately Long	Short	Extremely Short
(do not add to subtotal or total score)	0	1	2	3	4
7. INTERFERENCE FROM COMPULSION	0	1	2	3	4
8. DISTRESS FROM COMPULSIONS	0	1	2	3	4
9. RESISTANCE	Always resists 0	1	2	3	Completely yields 4
10. CONTROL OVER COMPULSIONS	Complete control 0	Much control 1	Moderate control 2	Little control 3	No control 4
COMPULSION SUBTOTAL (add items 6-10)					[      ]

19. RELIABILITY      EXCELLENT = 0      GOOD = 1      FAIR = 2      POOR = 3

EXCELLENT = no reason to suspect data unreliable; GOOD= factor(s) that may adversely affect reliability;  
 FAIR= factor(s) that definitely reduce reliability; POOR= very low reliability.



## Instructions to Clinicians Using the Obsession and Compulsion Log

The Obsession and Compulsion Log (OCL) provides two kinds of clinical information. First, it illustrates the nature of a patient's current obsessions and compulsions. Second, it is an index of clinical severity by providing information on the frequency of obsessions and the amount of time spent engaged in compulsive rituals.

Typically, patients are instructed to record all obsessions and compulsions that occur over the course of a single day. However, the time period within which obsessions and compulsions are recorded will vary according to the patient's condition. For example, when obsessions are relatively infrequent, the patient may need to complete the log for several days or longer to provide sufficient clinical data. On the other hand, patients with constant or very frequent obsessions may have difficulty completing the log for an entire 24-hour period. These patients can be instructed to record obsessions and compulsions for some designated time period less than 24 hours. When considering what time period to assign, remember the purpose of the OCL is to obtain a sample of obsessions and compulsions adequate enough to make clinical decisions.

NAME \_\_\_\_\_ DATE \_\_\_\_\_

DATE \_\_\_\_\_

Time	Obsession	Compulsion	Duration

Time

## Obsession

Compulsion

Duration

# DASS

Name:

Date:

Please read each statement and circle a number 0, 1, 2 or 3 that indicates how much the statement applied to you *over the past week*. There are no right or wrong answers. Do not spend too much time on any statement.

*The rating scale is as follows:*

- 0 Did not apply to me at all
- 1 Applied to me to some degree, or some of the time
- 2 Applied to me to a considerable degree, or a good part of time
- 3 Applied to me very much, or most of the time

1	I found myself getting upset by quite trivial things	0	1	2	3
2	I was aware of dryness of my mouth	0	1	2	3
3	I couldn't seem to experience any positive feeling at all	0	1	2	3
4	I experienced breathing difficulty (eg, excessively rapid breathing, breathlessness in the absence of physical exertion)	0	1	2	3
5	I just couldn't seem to get going	0	1	2	3
6	I tended to over-react to situations	0	1	2	3
7	I had a feeling of shakiness (eg, legs going to give way)	0	1	2	3
8	I found it difficult to relax	0	1	2	3
9	I found myself in situations that made me so anxious I was most relieved when they ended	0	1	2	3
10	I felt that I had nothing to look forward to	0	1	2	3
11	I found myself getting upset rather easily	0	1	2	3
12	I felt that I was using a lot of nervous energy	0	1	2	3
13	I felt sad and depressed	0	1	2	3
14	I found myself getting impatient when I was delayed in any way (eg, elevators, traffic lights, being kept waiting)	0	1	2	3
15	I had a feeling of faintness	0	1	2	3
16	I felt that I had lost interest in just about everything	0	1	2	3
17	I felt I wasn't worth much as a person	0	1	2	3
18	I felt that I was rather touchy	0	1	2	3
19	I perspired noticeably (eg, hands sweaty) in the absence of high temperatures or physical exertion	0	1	2	3
20	I felt scared without any good reason	0	1	2	3
21	I felt that life wasn't worthwhile	0	1	2	3

Please turn the page ➞

*Reminder of rating scale:*

- 0 Did not apply to me at all
- 1 Applied to me to some degree, or some of the time
- 2 Applied to me to a considerable degree, or a good part of time
- 3 Applied to me very much, or most of the time

22	I found it hard to wind down	0	1	2	3
23	I had difficulty in swallowing	0	1	2	3
24	I couldn't seem to get any enjoyment out of the things I did	0	1	2	3
25	I was aware of the action of my heart in the absence of physical exertion (eg, sense of heart rate increase, heart missing a beat)	0	1	2	3
26	I felt down-hearted and blue	0	1	2	3
27	I found that I was very irritable	0	1	2	3
28	I felt I was close to panic	0	1	2	3
29	I found it hard to calm down after something upset me	0	1	2	3
30	I feared that I would be "thrown" by some trivial but unfamiliar task	0	1	2	3
31	I was unable to become enthusiastic about anything	0	1	2	3
32	I found it difficult to tolerate interruptions to what I was doing	0	1	2	3
33	I was in a state of nervous tension	0	1	2	3
34	I felt I was pretty worthless	0	1	2	3
35	I was intolerant of anything that kept me from getting on with what I was doing	0	1	2	3
36	I felt terrified	0	1	2	3
37	I could see nothing in the future to be hopeful about	0	1	2	3
38	I felt that life was meaningless	0	1	2	3
39	I found myself getting agitated	0	1	2	3
40	I was worried about situations in which I might panic and make a fool of myself	0	1	2	3
41	I experienced trembling (eg, in the hands)	0	1	2	3
42	I found it difficult to work up the initiative to do things	0	1	2	3

# DASS

## Scoring Template

S  
A  
D  
A  
  
D  
S  
A  
S  
A  
  
D  
S  
S  
D  
S  
  
A  
D  
D  
S  
A  
  
A  
D

Apply template to both sides of sheet and sum scores for each scale.  
For short (21-item) version, multiply sum by 2.



# The Depression Anxiety Stress Scales (DASS): Normative data and latent structure in a large non-clinical sample

John R. Crawford\* and Julie D. Henry

Department of Psychology, King's College, University of Aberdeen, UK

**Objectives.** To provide UK normative data for the Depression Anxiety and Stress Scale (DASS) and test its convergent, discriminant and construct validity.

**Design.** Cross-sectional, correlational and confirmatory factor analysis (CFA).

**Methods.** The DASS was administered to a non-clinical sample, broadly representative of the general adult UK population ( $N = 1,771$ ) in terms of demographic variables. Competing models of the latent structure of the DASS were derived from theoretical and empirical sources and evaluated using confirmatory factor analysis. Correlational analysis was used to determine the influence of demographic variables on DASS scores. The convergent and discriminant validity of the measure was examined through correlating the measure with two other measures of depression and anxiety (the HADS and the sAD), and a measure of positive and negative affectivity (the PANAS).

**Results.** The best fitting model ( $CFI = .93$ ) of the latent structure of the DASS consisted of three correlated factors corresponding to the depression, anxiety and stress scales with correlated error permitted between items comprising the DASS subscales. Demographic variables had only very modest influences on DASS scores. The reliability of the DASS was excellent, and the measure possessed adequate convergent and discriminant validity.

**Conclusions.** The DASS is a reliable and valid measure of the constructs it was intended to assess. The utility of this measure for UK clinicians is enhanced by the provision of large sample normative data.

The Depression Anxiety Stress Scale (DASS) is a 42-item self-report measure of anxiety, depression and stress developed by Lovibond and Lovibond (1995) which is increasingly used in diverse settings. Its popularity is partly attributable to the fact

\*Requests for reprints should be addressed to John R. Crawford, Department of Psychology, King's College, University of Aberdeen AB24 3HN, UK (e-mail: [j.crawford@abdn.ac.uk](mailto:j.crawford@abdn.ac.uk)).



that, unlike many other self-report scales, the DASS is in the public domain (i.e. the measure can be used without incurring any charge). The DASS was originally intended to consist of only two subscales—one measuring anxiety, the other depression—each composed of items that were purportedly unique to either construct. Ambiguous items (i.e. items non-specifically related to depression and anxiety) were not included in the measure but were regarded as controls. This strategy was adopted because the authors' original intention was to develop measures that would maximally discriminate between depression and anxiety. However, during scale development it was revealed that the control items tended to form a third group, of items characterized by chronic non-specific arousal. More items were added to this group and the third scale, the stress scale, emerged. Lovibond and Lovibond maintain that, although this scale is related to the constructs of depression and anxiety, it nevertheless represents a coherent measure in its own right.

Whilst Lovibond and Lovibond's (1995) attempt to develop a measure that maximally discriminates between the constructs of depression and anxiety is not unique (Beck, Epstein, Brown, & Steer, 1988; Costello & Comrey, 1967), the strategy adopted for scale construction is. Conventionally, items are derived from pre-existing anxiety and depression scales, with factor analyses of clinical data used to identify those which measure different constructs. By contrast, Lovibond and Lovibond employed predominantly non-clinical samples for scale development on the basis that depression and anxiety represent dimensional, not categorical, constructs. Moreover, core symptoms of anxiety and depression which were unique to one but not both of the disorders were identified from the outset, and not on an *a posteriori* basis. Thus, unconventionally, the initial items selected were retained, with new items compatible with the emerging factor definitions successively added.

Preliminary evidence has been presented, which suggests that the DASS does possess adequate convergent and discriminant validity (Lovibond & Lovibond, 1995). A large student sample ( $N = 717$ ) was administered the Beck Depression Inventory (BDI; Beck, Ward, Mendelsohn, Mock, & Erbaugh, 1961), the Beck Anxiety Inventory (BAI; Beck *et al.*, 1988) and the DASS. The BAI and DASS anxiety scale were highly correlated ( $r = .81$ ), as were the BDI and DASS depression scale ( $r = .74$ ). However, between-construct correlations were substantially lower ( $r = .54$  for DASS depression and BAI;  $r = .58$  for DASS anxiety and BDI). Moreover, Antony, Bieling, Cox, Enns, and Swinson (1998) found a similar pattern of correlations in a clinical sample.

To assess the DASS's psychometric properties, Lovibond and Lovibond (1995) administered the measure to a large non-clinical sample ( $N = 2,914$ ). It was found that reliability, assessed using Cronbach's alpha, was acceptable for the depression, anxiety and stress scales (.91, .84 and .90, respectively). These values are similar to those obtained from clinical populations (Antony *et al.*, 1998; Brown, Chorpita, Korotitsch, & Barlow, 1997).

At present, interpretation of the DASS is based primarily on the use of cut-off scores. Lovibond and Lovibond (1995) presented severity ratings from 'normal' to 'extremely severe' on the basis of percentile scores, with 0–78 classified as 'normal', 78–87 as 'mild', 87–95 as 'moderate', 95–98 as 'severe', and 98–100 as 'extremely severe'. However, these original norms were based predominantly on students. This means that the generalizability of their results to the normal population is uncertain. Moreover, although 1,307 of the participants in this study were non-students, no information was presented regarding whether they were broadly representative of the general

population; all that was stated was that they were 'white and blue collar workers' (Lovibond & Lovibond, 1995, p. 9).

Relatedly, the influence of demographic characteristics on DASS scores has gone largely uninvestigated. In development of the DASS, this analysis was restricted to gender and age. Although the test authors did not state explicitly whether age and/or gender yielded a significant effect, '... there was a trend towards higher scores in the youngest and oldest age brackets' (Lovibond & Lovibond, 1995, p. 28). However, Andrew, Baker, Kneebone, and Knight (2000) found that in a sample of elderly community volunteers ( $N = 53$ ), scores on all three DASS subscales were almost half those reported by Lovibond and Lovibond. It is possible that this discrepancy is attributable to idiosyncrasies in one or both of these samples or the influence of potential mediating factors such as years of education or occupation. Yet no study to date has assessed the influence of either of these latter variables. The relationships between demographic variables and DASS scores in the general population are of interest in their own right, but investigation of these relationships would also serve the very practical purpose of identifying whether normative data should be stratified.

If the use of the DASS in research and clinical practice is to be optimal, then it is also necessary to delineate the underlying structure of the instrument. This is particularly important given that Lovibond and Lovibond (1995) found through empirical analyses that, in both clinical and non-clinical samples, symptoms conventionally regarded as core to the syndrome of depression (American Psychiatric Association, 1994) were actually extremely weak markers of this construct. Specifically, items pertaining to changes in appetite, sleep disturbance, guilt, tiredness, concentration loss, indecision, agitation, loss of libido, diurnal variation in mood, restlessness, irritability and crying were excluded from the measure.

Moreover, the legitimacy of the stress scale as an independent measure must be assessed. In an influential series of papers, Clark and Watson (Clark & Watson, 1991a, 1991b; Watson, Clark, & Tellegen, 1988) have argued that anxiety and depression have an important shared component which they call 'negative affectivity' (NA). NA is a dispositional dimension, with high NA reflecting the experience of subjective distress and unpleasurable engagement, manifested in a variety of emotional states such as guilt, anger and nervousness, and low NA represented by an absence of these feelings (Watson & Clark, 1984). Studies have supported the existence of a dominant NA dimension (Watson & Clark, 1984; Watson & Tellegen, 1985) and provide evidence that it is highly related to the symptoms of both anxiety and depression (Brown *et al.*, 1997; Watson, Clark, Weber *et al.*, 1995; Watson, Weber *et al.*, 1995). Thus, there are strong theoretical grounds for suggesting that the stress scale is simply a measure of NA, particularly given that this scale actually originated from items believed to relate to *both* dimensions.

To date, four studies have directly tested the construct validity of the DASS (Antony *et al.*, 1998; Brown *et al.*, 1997; Clara, Cox, & Enns, 2001; Lovibond & Lovibond, 1995). Lovibond and Lovibond (1995) conducted a principal-components analysis in a student sample ( $N = 717$ ) which revealed that the first three factors accounted for a high proportion of the variance. Furthermore, all items loaded on their designated factor except for anxiety item 30 ('I feared that I would be "thrown" by some trivial but unfamiliar task') which loaded on the stress factor. In the same sample, a confirmatory factor analysis (CFA) was then used to quantitatively compare the fit of a single-factor model, a two-factor model (in which depression was one factor, and anxiety and stress were collapsed into another) and a three-factor model corresponding to the three DASS

scales. The three-factor model was found to represent the optimal fit, and a significantly better fit than the two-factor model.

Analogous findings have been reported in two independent clinical samples. Brown *et al.* (1997) conducted an exploratory factor analysis (EFA) with varimax rotation using data derived from a sample ( $N = 437$ ) of patients suffering from a range of affective disorders. A three-factor solution emerged, reproducing Lovibond and Lovibond's (1995) hypothesized structure. The only discrepancies were that anxiety item 9 ('I found myself in situations which made me so anxious that I was most relieved when they ended') and stress item 33 ('I was in a state of nervous tension')<sup>1</sup> double loaded, and anxiety item 30 failed to load strongly on any factor. Brown *et al.* then administered the instrument to an independent clinical sample ( $N = 241$ ) and employed CFA to test the fit of four models. The first three models corresponded exactly to those tested by Lovibond and Lovibond. In addition, a model revised according to the results of the EFA conducted with Brown *et al.*'s first sample was also tested. The results revealed that the revised model represented the optimal fit, and a significantly better fit than the model corresponding to Lovibond and Lovibond's original specifications.

Finally, both Clara *et al.* (2001) and Antony *et al.* (1998) identified three factor solutions in clinical samples ( $N = 258$  and  $N = 439$ , using CFA and EFA respectively). Antony *et al.* (1998), however, again noted discrepancies; stress items 22 ('I found it hard to wind down') and 33 double loaded on anxiety, and anxiety items 9 and 30 double loaded on stress. Thus, whilst these studies suggest that there is a slight degree of misspecification, they have consistently supported the validity of a three-factor structure corresponding to the dimensions of anxiety, depression and stress. To date, though, no study has tested the construct validity of the DASS in a sample drawn from the general adult population.

The aims of the present study were:

- (1) to investigate the influence of demographic variables on DASS scores in the general adult UK population;
- (2) to provide UK normative data for the DASS in the form of tables for converting raw scores to percentiles;
- (3) to evaluate competing models of the latent structure of the DASS using CFA (details of the parameterization of the models, and the theoretical, methodological and empirical considerations that guided their selection, are presented in the methods section);
- (4) to obtain estimates of the reliability of the DASS; and
- (5) to test the convergent and discriminant validity of the DASS.

## Method

### Participants

Complete DASS data were collected from 1,771 members of the general adult population (females = 965, males = 806). Participants were recruited from a wide variety of sources including commercial and public service organizations, community centres and recreational clubs. The mean age of the sample was 40.9 ( $SD = 15.9$ ) with a range of 15–91 years. The mean years of education was 13.8 ( $SD = 3.1$ ).

---

<sup>1</sup> Brown *et al.* (1997) refer to item 33 as item 34.

### **Materials and procedure**

Each potential participant received an introductory letter, a DASS form, and a form for recording demographic variables. A subset of participants also received and completed two additional self-report measures of depression and anxiety, as well as a measure of positive and negative affect. These were the Hospital Anxiety and Depression Scale (HADS,  $N = 1512$ ; Zigmond & Snaith, 1983), the Personal Disturbance scale (sAD,  $N = 733$ ; Bedford & Foulds, 1978), and the Positive and Negative Affect Schedule (PANAS,  $N = 740$ ; Watson *et al.*, 1988). Participants sealed the completed forms in an envelope, and these were either collected by the researcher or returned by mail. The refusal rate was approximately 18% (participants who failed to return forms or returned entirely blank forms were also treated as refusals). In addition, of the 1,786 completed forms, 15 contained either some missing data or contained equivocal responses; these forms were discarded.

Each participant's occupation was coded using the Office of Population Censuses and Surveys (1990) *Classification of occupations*. Retired participants, and those describing themselves as househusbands/housewives, were coded by their previous occupations, as were those currently unemployed. Those who had never worked were coded as 5 (i.e. unskilled).

The percentage of participants in the occupational codes of professional (1), intermediate (2), skilled (3), semi-skilled (4) and unskilled (5) was 11, 38, 34, 9 and 8, respectively. The corresponding percentage for each code in the general adult population census is 7, 32, 42, 14 and 5, respectively. Thus, whilst there was a broad spread, there was a slight overrepresentation of professional occupations, and a slight underrepresentation of skilled and semi-skilled occupations. The percentage of participants in each of four age bands (18–29, 30–44, 45–59, 60+) was 30, 31, 26 and 14. The corresponding percentage for each age band in the general adult population census is 27, 25, 22 and 26, respectively. Again it can be seen that there was a broad spread, although there was a relative underrepresentation of individuals in the oldest age group.

#### *The Hospital Anxiety and Depression Scale (HADS)*

The HADS was developed by Zigmond and Snaith (1983) to provide a brief means of identifying and measuring severity of depression and anxiety in non-psychiatric clinical environments. It consists of 14 items, seven of which measure depression, the other seven anxiety. The respondent is asked to underline the reply which most closely matches how they have felt during the past week.

#### *The Personal Disturbance Scale (sAD)*

The sAD is a brief (14-item) self-report measure derived from the Delusions-Symptoms States Inventory (DSSI; Bedford & Foulds, 1978), and consists of seven anxiety and seven depression items.

#### *The Positive and Negative Affect Schedule (PANAS)*

The PANAS is a brief (20-item) self-report measure of positive affect and negative affect developed by Watson *et al.* (1988). It is claimed that the PANAS provides independent (i.e. orthogonal) measures of these constructs. The 'past week' time format was adopted.

### Statistical analysis

Basic statistical analysis was conducted using SPSS Version 8. Confidence limits on Cronbach's alpha were derived from Feldt's (1965) formulae.

CFA (robust maximum likelihood) was performed on the variance-covariance matrix of the DASS items using EQS for Windows Version 5 (Bentler, 1995). The fit of CFA models was assessed using the Satorra-Bentler scaled chi square statistic (S-B  $\chi^2$ ), the average off-diagonal standardized residual (AODSR), the Comparative Fit Index (CFI), the Robust Comparative Fit Index (RCFI) and the Root Mean Squared Error of Approximation (RMSEA). Off-diagonal standardized residuals reflect the extent to which covariances between observed variables have not been accounted for by the models under consideration. Values for the CFI and RCFI can range from zero to unity; these indices express the fit of a model relative to what is termed the 'null model' (the null model posits no relationship between any of the manifest variables). There is general agreement that a model with a CFI of less than 0.95 should not be viewed as providing a satisfactory fit to the data (Hu & Bentler, 1999). The RMSEA has been included as this fit index explicitly penalizes models which are not parsimonious.

A model is considered to be nested within another model if it differs only in imposing additional constraints on the relationships between variables specified in the initial model. The difference between chi square for nested models is itself distributed as chi square with  $k$  degrees of freedom where  $k$  equals the degrees of freedom for the more constrained model minus the degrees of freedom for the less constrained model. This means that it is possible to test directly whether more constrained models have a significantly poorer fit than less constrained models; this feature of CFA is one of its major advantages over EFA. In the present case there is a slight complication because the S-B  $\chi^2$  is used as an index of fit rather than the standard chi-square statistic (the Satorra-Bentler statistic is recommended when the raw data are skewed). The *difference* between S-B  $\chi^2$  for nested models is typically not distributed as chi square. However, Satorra and Bentler (2001) have recently developed a scaled difference chi-square test statistic that can be used to compare S-B  $\chi^2$  from nested models. This statistic is used in the present study.<sup>2</sup>

### Parameterization of competing models of the DASS

The first model (Model 1a) to be evaluated was a single-factor model; this model expressed the hypothesis that the variance in the DASS can be partitioned into one general factor plus error variance associated with each individual item. It is standard practice to test the fit of a one-factor model because it is the most parsimonious of all possible models. A further model was tested (Model 1b) in which again all items were presumed to load upon only one general factor. However, as can be seen in Table 6, items in each of the DASS scales are grouped into categories hypothesized to measure the same subcomponents of the relevant construct. In Model 1b, items from the same content categories were permitted to covary. No study to date has tested a model parameterized to allow for such correlated error.

Models 2a-2c expressed variants on the hypothesis that the DASS measures two factors, anxiety and depression. For all three models the items in the stress and anxiety scale were collapsed into one factor to test the hypothesis that the stress scale does not represent an independent construct but, rather, simply measures anxiety. In Model 2a

<sup>2</sup> In the course of analysing the present data we wrote a computer program (for PCs) that carries out this test. The program can be downloaded from [www.psyc.abdn.ac.uk/homedir/jcrawford/sbdiff.htm](http://www.psyc.abdn.ac.uk/homedir/jcrawford/sbdiff.htm)

these two factors were constrained to be orthogonal and in Model 2b, permitted to correlate. Model 2b was then retested, but additionally permitted correlated error between items from the same content categories (Model 2c).

Models 3a–3d tested Lovibond and Lovibond's (1995) three-factor structure, specifying the dimensions of anxiety, depression and stress. In Model 3a, the three factors were constrained to be orthogonal, with Model 3b permitting the factors to correlate in accordance with Lovibond and Lovibond's original specifications. Model 3c represented a test of the model which Brown *et al.* (1997) derived through an EFA in a clinical sample, and which represented the optimal fit of four CFA models tested in an independent clinical sample. The model was parameterized according to Lovibond and Lovibond's original specifications, except that some items were permitted to load on more than one factor. Specifically, stress item 33 also loaded on anxiety, anxiety item 9 on stress, and anxiety item 30 on all three factors. Finally, Model 3c was retested, but additionally permitted correlated error (Model 3d).

## Results

### *Influence of demographic variables on DASS scores*

As the DASS scales had a high positive skew, analysis of their relationships with demographic variables (i.e. *t*-tests and correlations) was performed on the logarithm of their scores. Independent samples *t*-tests revealed that females obtained significantly higher scores than males on the anxiety scale ( $M = 4.0$ ,  $SD = 6.17$  [females];  $M = 3.0$ ,  $SD = 4.23$  [males];  $t = -2.29$ ,  $p < .05$ ), depression scale ( $M = 6.1$ ,  $SD = 8.14$  [females];  $M = 4.9$ ,  $SD = 6.55$  [males];  $t = -2.68$ ,  $p < .01$ ), and total of the three scales ( $M = 19.9$ ,  $SD = 20.82$  [females];  $M = 16.6$ ,  $SD = 15.95$  [males];  $t = -2.20$ ,  $p < .05$ ). The difference between males and females on the stress scale did not achieve statistical significance ( $M = 9.8$ ,  $SD = 8.56$  [males];  $M = 8.7$ ,  $SD = 7.35$  [females];  $t = -1.802$ ,  $p > .05$ ).

**Table 1.** Correlations between demographic variables and DASS scores

Demographic variable	DASS			
	Anxiety	Depression	Stress	Total
Age	-.036	-.109**	-.183**	-.147**
Occupational code	.066**	.018	-.039	.005
Years of education	-.033	-.008	.086**	-.054*
Gender	.054*	.064**	.043	.052*

\* Correlation significant at .05 level (two-tailed); \*\* correlation significant at .01 level (two-tailed).

The influence of the remaining demographic variables (age, years of education and occupational code) on the DASS anxiety, depression, stress and total scales was tested through correlational analyses, the results of which are presented in Table 1. The point-biserial correlations between gender and the DASS scale scores are also presented in this table as an index of effect size (males were coded as 0, females as 1, so a positive

correlation represents a higher score in females). It can be seen from Table 1 that the influence of all demographic variables on DASS scores is very modest.

### Summary statistics and normative data for the DASS

The means, medians, *SDs* and ranges for each of the three DASS scales are presented in Table 2 for the total sample. Additionally, for each subscale the percentage of participants falling into each of the five categories (normal, mild, moderate, severe and extremely severe) created by the use of Lovibond and Lovibond's (1995) cut-off scores is presented. However, these cut-offs have been presented purely for comparative purposes, and it is important to reiterate that DASS scores should be regarded as providing an individual's score on an underlying dimension.

**Table 2.** Summary statistics for DASS

	Median	<i>M</i>	<i>SD</i>	Range	Percentage in each DASS category				
					Normal (0–78 <sup>a</sup> )	Mild (78–87)	Moderate (87–95)	Severe (95–98)	Extremely severe (98–100)
Total sample ( <i>N</i> = 1771)									
Anxiety	2	3.56	5.39	0–40	94.4	2.0	3.8	2.0	3.2
Depression	3	5.55	7.48	0–42	81.7	6.2	6.3	2.9	2.9
Stress	8	9.27	8.04	0–42	80.2	8.4	5.9	3.5	2.0
Total	13	18.38	18.82	0–121					

<sup>a</sup> Lovibond and Lovibond's (1995) percentile cut-offs corresponding to each DASS category.

Visual inspection of the distribution of raw scores on the four scales revealed that, as is to be expected in a sample drawn from the general adult population, they were positively skewed, particularly the anxiety scale. Kolmogorov–Smirnov tests confirmed that the distributions deviated highly significantly from a normal distribution (*Z* ranged from 5.24 to 10.70, all *ps* < .001).

Given the positive skew, use of the means and *SDs* from a normative sample is not useful when interpreting an individual's score. Therefore, Table 3 was constructed for conversion of raw scores on each of the DASS scales to percentiles.

### Testing competing confirmatory factor analytic models of the DASS

The fit statistics for the CFA models are presented in Table 4. It can be seen that the general factor model (Model 1a) had a very poor fit; the  $\chi^2$  is large, and the fit indices are low. However, all items loaded highly on this factor, evidence that there is substantial common variance among the items. Permitting correlated error (Model 1b) led to an improved, but still badly fitting, model. The two-factor models also had a poor fit, although the correlated factors models (Models 2b and 2c) were better than their more constrained counterpart (Model 2a). Again, correlated error led to an improvement in fit (Model 2c having higher fit indices and a lower  $\chi^2$  than Model 2b).

Model 3a tested Lovibond and Lovibond's (1995) three-factor structure but specified orthogonal constructs. This was associated with low fit indices and a very high  $\chi^2$ . Although permitting correlated factors in Model 3b improved the model's fit, it was still

**Table 3.** Raw scores on the DASS converted to percentiles

Percentile	Raw scores				Percentile
	Depression	Anxiety	Stress	Total	
5	0	0	0	1	5
10	0	0	1	2	10
15	0	0	2	3	15
20	0	0	3	5	20
25	1	0	3	6	25
30	1	0	4	7	30
35	1	1	5	8	35
40	2	1	6	10	40
45	2	1	7	12	45
50	3	2	8	13	50
55	3	2	8	15	55
60	4	3	9	17	60
65	5	3	10	19	65
70	6	4	12	22	70
75	7	4	13	24	75
76	8	5	13	24	76
77	8	5	13	25	77
78	8	5	14	26	78
79	9	5	14	27	79
80	9	6	14	28	80
81	9	6	15	28	81
82	10	6	15	29	82
83	10	6	16	30	83
84	11	7	16	31	84
85	11	7	17	32	85
86	12	7	17	34	86
87	13	8	18	35	87
88	14	8	18	36	88
89	14	8	19	39	89
90	15	9	20	40	90
91	16	10	21	42	91
92	17	11	22	46	92
93	18	12	23	48	93
94	20	13	25	54	94
95	22	15	26	60	95
96	24	17	28	64	96
97	27	20	30	72	97
98	31	22	34	79	98
99	36	26	37	91	99

poor. However, for both models, all items loaded highly on the appropriate construct. Model 3c represented a revised version of Lovibond and Lovibond's model based on the empirical findings of Brown *et al.* (1997) and represented a superior fit. As with Brown *et al.*'s study, items 9 and 33 loaded equivalently on the anxiety and stress factors (.36 vs. .36; .41 vs. .40, respectively), and item 30 loaded weakly on all three factors (ranging from .12 to .35). Again, none of the fit indices was acceptable. Model 3d was identical to



**Table 4.** Fit indices for CFA models of DASS

Model	S-B $\chi^2$	$\chi^2$ <sup>a</sup>	df	AODSR	CFI	RCFI	RMSEA
Single factor							
1a. Single factor	7,259.3	14,144.5	819	.0560	.726	.542	.096
1b. Single factor with correlated error	3,986.4	7,616.1	779	.0475	.860	.772	.070
Anxiety and depression as...							
2a. independent factors	6,172.2	11,902.2	819	.2063	.773	.619	.087
2b. correlated factors	5,421.9	10,341.7	818	.0459	.805	.673	.081
2c. correlated factors with correlated error	2,965.0	5,607.6	778	.0385	.901	.844	.059
Lovibond & Lovibond's model with...							
3a. independent factors	5,661.8	10,945.0	819	.2662	.792	.656	.084
3b. correlated factors	4,298.2	8,148.0	816	.0422	.850	.752	.071
3c. correlated factors, revised	4,059.5	7,656.9	812	.0377	.860	.769	.069
3d. correlated factors, revised, and correlated error	2,347.8	4,403.2	772	.0322	.925	.888	.052

<sup>a</sup> The Satorra-Bentler scaled chi square statistic (S-B  $\chi^2$ ) was used to evaluate model fit. However, the normal chi square is also required when testing for a difference between the S-B  $\chi^2$  statistic obtained from nested models; hence we present both statistics in this table.

**Table 5.** Results of testing for differences between nested CFA models of DASS

Comparison		$\Delta$ statistics		
More constrained	Less constrained	$\Delta$ S-B $\chi^2$	df	<i>p</i>
Model 1a	Model 1b	3,272.9	40	<.001
Model 2a	Model 2b	750.3	1	<.001
Model 1a	Model 2b	1,837.4	1	<.001
Model 2b	Model 2c	2,457.0	40	<.001
Model 1b	Model 2c	1,021.4	1	<.001
Model 3a	Model 3b	1,363.6	3	<.001
Model 1a	Model 3b	2,961.1	3	<.001
Model 2b	Model 3b	1,123.7	2	<.001
Model 3b	Model 3c	238.7	4	<.001
Model 3c	Model 3d	1,711.7	40	<.001
Model 1b	Model 3d	1,638.6	7	<.001
Model 2c	Model 3d	617.2	6	<.001

Model 3c but additionally permitted correlated error. This model was associated with the optimal fit according to all criteria, with high fit indices and a  $\chi^2$  value that, although statistically significant,<sup>3</sup> was substantially lower than that for the other models tested.

The fit of the correlated factors models is markedly superior to their independent factors counterparts. As noted, inferential statistics can be applied to compare nested models. Models 2a and 3a are nested within Models 2b and 3b respectively in that they differ only by the imposition of the constraint that the factors are independent. The results from chi square difference tests used to compare these nested models are presented in Table 5. It can be seen that the correlated factors models had a significantly better fit ( $p < .001$ ) than their independent factors counterparts, demonstrating that the conception of independence between the scales is untenable. This is underlined by the correlations between the three *factors* in Models 3b–3d. For the optimal Model, 3d, the correlations were depression-anxiety ( $r = .75$ ), stress-depression ( $r = .77$ ) and stress-anxiety ( $r = .74$ ). These correlations are higher than the respective correlations between the *scales*: depression-anxiety ( $r = .70$ ), stress-depression ( $r = .72$ ) and stress-anxiety ( $r = .70$ )—although these latter correlations are themselves substantial. This is because the factors in the CFA models are measured without error, whereas the correlation between the scales is attenuated by measurement error and the unique variance associated with each item.

Although it may appear initially that the general factor model is very different from the correlated factors models, it is also nested within these models. Models 2b and 3b can be rendered equivalent to a single factor simply by constraining the correlation between factors to unity (i.e.  $r = 1.0$ ). The chi square difference tests comparing Model 1 with Models 2b and 3b were both highly significant, demonstrating that it is also untenable to view the DASS as measuring only a single factor of negative affectivity or general psychological distress.

Allowing for correlated error between the items also resulted in a significant

<sup>3</sup> When dealing with large sample sizes and a large number of items it is unusual to obtain non-significant  $\chi^2$  values for CFA models of self-report data (Byrne, 1994).

improvement in the fit of Models 1b, 2c and 3d compared with their more constrained counterparts, Models 1a, 2b and 3c, respectively. Moreover, the addition of the double loadings identified by Brown *et al.* (1997) led to improvement, with Model 3c a significantly better fit than Model 3b ( $p < .001$ ).

### **Evaluation of the optimal model**

As shown in Table 6, all items in Model 3d loaded  $\geq .47$  on the specific factor they were intended to represent, with the exception of the three 'weak' items identified in earlier factor analyses (items 9, 30 and 33). Cross-validating Brown *et al.*'s (1997) clinical study, anxiety item 9 and stress item 33 loaded identically on the anxiety and stress factors (item 9 loaded .36 on both factors; and item 33 loaded .40 on each construct). Item 30 loaded only weakly on all three factors (.13, .36 and .23 on depression, anxiety and stress, respectively). Although allowing correlated error between items of related subscales led to a significant improvement in fit, the item-specific correlations revealed that not all of the subsets appeared to be related in the manner hypothesized. That is, although the majority were positively related, some correlations were negative, albeit modestly so.

A schematic representation of the structure for the optimal Model (3d) is presented as Figure 1 (the associated factor loadings are presented in Table 6). By convention, latent factors are represented by large ovals or circles, the error variances as smaller ovals or circles (as they are also latent variables) and manifest (i.e. observed) variables as rectangles or squares. Single-headed arrows connecting variables represent a causal path. Double-headed arrows represent covariance or correlation between variables but do not imply causality.

### **Reliabilities of the DASS**

The reliabilities (internal consistencies) of the DASS anxiety, depression, stress and total score were estimated using Cronbach's alpha. Alpha was .897 (95% CI = .890-.904) for the anxiety scale, .947 (95% CI = .943-.951) for the depression scale, .933 (95% CI = .928-.937) for the stress scale, and .966 (95% CI = .964-.968) for the total score.


### **Convergent and discriminant validity of the DASS**

To examine the convergent and discriminant validity of the DASS, Pearson product-moment correlations were calculated between each of the DASS scales and the sAD, HADS and PANAS scales. These correlations are presented in Table 7. With respect to convergent validity, the DASS depression scale correlated highly with sAD depression (.78). William's (1959) test revealed that this correlation was higher than that between sAD depression and HADS depression (.58;  $t = 9.10$ ,  $p < .001$ ). Similarly, the correlation between DASS depression and HADS depression (.66) was significantly higher than the HADS-sAD correlation ( $t = 4.19$ ,  $p < .001$ ). DASS anxiety scores also exhibited a high convergent validity. The correlation between DASS anxiety and sAD anxiety (.72) was significantly higher than that between the sAD and HADS anxiety scales (.67;  $t = 2.40$ ,  $p < .05$ ). However, although the correlation between DASS anxiety and HADS anxiety was substantial and highly significant (.62,  $p < .001$ ), it was lower than the aforementioned correlation between HADS anxiety and sAD anxiety ( $t = 2.41$ ,  $p < .05$ ).

**Table 6.** DASS items with factor loadings from confirmatory factor analysis (Model 3d)

Scale/item summary	Factor			
	Subscale	Depression	Anxiety	Stress
<b>Depression</b>				
26 Downhearted & blue	DYS	.77		
13 Sad & depressed	DYS	.78		
37 Nothing future hopeful	HLNS	.82		
10 Nothing to look forward to	HLNS	.81		
38 Life meaningless	DoL	.78		
21 Life not worthwhile	DoL	.79		
34 Felt worthless	S-Dep	.80		
17 Not worth much as person	S-Dep	.77		
16 Lost interest in everything	Lol/I	.81		
31 Unable to become enthusiastic	Lol/I	.78		
3 Couldn't experience positive	ANH	.71		
24 Couldn't get enjoyment	ANH	.75		
5 Couldn't get going	INRT	.53		
42 Difficult to work up initiative	INRT	.64		
<b>Anxiety</b>				
25 Aware of action of heart	AutAr		.62	
19 Perspired noticeably	AutAr		.60	
2 Dryness of mouth	AutAr		.47	
4 Breathing difficulty	AutAr		.50	
23 Difficulty swallowing	AutAr		.57	
7 Shakiness	SkME		.63	
41 Trembling	SkME		.62	
40 Worried about situations/panic	SitAnx		.62	
9 Situations made anxious	SitAnx		.36	.36
30 Feared would be 'thrown'	SitAnx	.13	.36	.23
28 Felt close to panic	SubAA		.80	
36 Terrified	SubAA		.70	
20 Scared for no good reason	SubAA		.74	
15 Feeling faint	SubAA		.58	
<b>Stress</b>				
22 Hard to wind down	DRel			.69
29 Hard to calm down	DRel			.79
8 Difficult to relax	DRel			.68
12 Using nervous energy	NerAr			.67
33 State of nervous tension	NerAr		.40	.40
11 Upset easily	EU/A			.79
1 Upset by trivial things	EU/A			.69
39 Agitated	EU/A			.78
6 Overreact to situations	I/OR			.72
27 Irritable	I/OR			.77
18 Touchy	I/OR			.76
35 Intolerant kept from getting on	IMPT			.62
14 Impatient when delayed	IMPT			.53
32 Difficulty tolerating interruptns	IMPT			.63

*Note.* DYS = dysphoria; HLNS = hopelessness; DoL = devaluation of life; S-Dep = self-deprecation; Lol/I = lack of interest/involvement; ANH = anhedonia; INRT = inertia; AutAr = autonomic arousal; SkME = skeletal musculature effects; SitAnx = situational anxiety; SubAA = subjective anxious affect; DRel = difficulty relaxing; NerAr = nervous arousal; EU/A = easily upset/agitated; I/OR = irritable/over-reactive; IMPT = impatient.



**Figure 1.** Graphical representation of a correlated three-factor model of the DASS (Model 3d); cross-loadings have been omitted in the interests of clarity.

**Table 7.** Correlations between the DASS, sAD, HADS and PANAS

	DASS depression	DASS anxiety	DASS stress	sAD depression	sAD anxiety	HADS depression	HADS anxiety	PANAS PA	PANAS NA
DASS depression	–	–	–	–	–	–	–		
DASS anxiety	.70 (1771)	–	–	–	–	–	–		
DASS stress	.72 (1771)	.71 (1771)	–	–	–	–	–		
sAD depression	.78 (733)	.56 (733)	.56 (733)	–	–	–	–		
sAD anxiety	.62 (733)	.72 (733)	.67 (733)	.70 (758)	–	–	–		
HADS depression	.66 (1512)	.49 (1512)	.54 (1512)	.58 (746)	.52 (746)	–	–		
HADS anxiety	.59 (1512)	.62 (1512)	.71 (1512)	.54 (746)	.67 (746)	.53 (1792)	–		
PANAS PA	–.48 (740)	–.29 (740)	–.31 (740)	– (0)	– (0)	–.52 (989)	–.31 (989)	–	
PANAS NA	.60 (740)	.60 (740)	.67 (740)	– (0)	– (0)	.44 (989)	.65 (989)	–.24 (1003)	–

Note. N for each correlation in parentheses.

In common with all other self-report scales of anxiety and depression, the discriminant validity of the DASS was less impressive: the between-construct correlations (i.e. DASS anxiety with HADS depression, etc.) were all highly significant (see Table 7). However, Williams' tests revealed that when the DASS scales were paired with their opposites from the other scales, all these latter (between-construct) correlations were significantly lower ( $p < .05$  or beyond) than the corresponding within-construct correlations referred to above.

The correlations between PA and NA with the DASS scales are of particular interest, especially the correlations between PA and the depression scale, and NA and the stress scale. The depression scale's correlation with PA was highly significant and negative in sign ( $-.48$ ); thus scoring high on depression was associated with low levels of PA. Using Meng, Rosenthal, and Rubin's (1992) method of comparing sets of non-independent correlations, this correlation was significantly higher than the correlations between PA and the other two DASS scales ( $-.29$  for anxiety and  $-.31$  for stress;  $z = 8.36$ ,  $p < .001$ ). The correlation between the stress scale and NA ( $.67$ ) was significantly higher than the correlation of NA with the other two DASS scales ( $.60$  for both anxiety and depression;  $z = 3.64$ ,  $p < .001$ ).

## Discussion

### *Influence of demographic variables*

One basic aim of the present study was to examine the influence of demographic variables on DASS scores. Although nine out of the 16 relationships examined proved significant, the size of the effects was very modest. The percentage of variance explained ranged from a low of 0.003% (occupational code and total score) to 3.35% (age and stress). Thus, for practical purposes, the influence of gender, occupation, education and age on DASS scores can be ignored; the significant effects result from the high statistical power conferred by a large sample size. This simplifies interpretation of DASS scores, as these variables do not need to be taken into consideration.

The effects of gender on DASS scores were very modest; the largest effect was on the depression scale, but even here gender only accounted for 0.41% of the variance in scores. This result is surprising given that epidemiological studies generally report a higher incidence of anxiety and depression in females (Horwath & Weissman, 1995; Meltzer, Gill, Petticrew, & Hinds, 1995). It is not clear why substantial gender effects did not emerge in the present study, but this finding is consistent with Lovibond and Lovibond's (1995) study in which gender effects were also very modest. The explanation may lie in the combination of two factors. First, epidemiological studies are concerned with caseness, in other words only with the number of individuals that meet clinical criteria, rather than measuring milder manifestations of psychological distress. Second, the DASS intentionally omits many of the symptoms that form part of traditional psychiatric criteria (Lovibond & Lovibond, 1995).

### *Normative data*

Despite the widespread use of the DASS in the English-speaking world, adequate normative data for the English language version do not appear to have been presented previously. Instead, interpretation of the DASS has been based primarily on norms

derived from a sample predominantly composed of students (Lovibond & Lovibond, 1995). The current study usefully complements this by providing normative data derived from a sample known to be broadly representative of the general adult population.

The tabulation method in Table 3 was adopted to permit conversion of raw scores to percentiles for all three scales and the total scale using the same table. Because of this, and because of the granularity of raw scores, it can be seen that, in a few cases, a given raw score can correspond to more than one percentile (e.g. for the stress scale a raw score of 8 spans the 50th to 55th percentiles). When this occurs the user should take the highest percentile. Should clinicians or researchers prefer to express an individual's standing on the DASS as a normalized  $z$  score or  $T$  score, it would be a relatively simple task to derive these from the percentile tables. For example, a raw score of 22 on the DASS depression scale would convert to a  $T$  score of 66 or a  $z$  score of 1.64, given that this raw score corresponds to the 95th percentile.

The only previous normative data for the DASS comes from Lovibond and Lovibond's (1995) Australian sample. The mean score in the present sample for depression was 5.55 ( $SD = 7.48$ ), for anxiety 3.56 ( $SD = 5.39$ ) and for stress 9.27 ( $SD = 8.04$ ). These means are slightly lower than the norms presented by Lovibond and Lovibond: depression = 6.34 ( $SD = 6.97$ ); anxiety = 4.70 ( $SD = 4.91$ ) and stress = 10.11 ( $SD = 7.91$ ). The minor differences may be because Lovibond and Lovibond's data were derived from a sample predominantly composed of students; there is evidence of elevated rates of psychological disturbance in student populations (Boyle, 1985; Gotlib, 1984).

### **Competing models of the structure of the DASS**

CFA was used to test competing models of the latent structure of the DASS. From the fit statistics in Table 4, it is clear that the hypothesis that the DASS measures a single factor (Models 1a and 1b) is untenable. Analogously, all the two-factor models (Models 2a–2c) were associated with poor fits, although permitting correlated factors (Models 2b and 2c) and correlated error (Model 2c) each led to an improvement.

Model 3a represented a test of Lovibond and Lovibond's (1995) three-factor structure, but specified orthogonal factors. This model, as well as a model in which correlated factors were permitted (Model 3b), represented poor fits, with large  $\chi^2$  and low fit indices. However, both had a significantly better fit than their more constrained one-factor counterparts (Models 1a and 1b, respectively) and Model 3b's two-factor counterpart (Model 2b). This indicates that it is untenable to view the DASS as measuring only one or two dimensions; the stress scale represents a legitimate construct in its own right. Model 3c, identical to Model 3b except for Brown *et al.*'s (1997) empirically derived revisions, represented a significantly better fit than Model 3b. However, additionally permitting correlated error between related subscales resulted in the optimal fitting model, as reflected by all criteria, even though some of the subscales appear to consist of items that are heterogeneous in content.

The conclusion from the CFA modelling, therefore, is that, consistent with previous empirical findings, the depression, anxiety and stress scales do represent legitimate constructs in their own right. Moreover, the current study supports Brown *et al.*'s (1997) findings that minor adjustments are required to optimize fit.



### **Reliabilities**

The reliabilities of the DASS scales, as measured by Cronbach's alpha, were .90 for anxiety, .95 for depression, .93 for stress and .97 for the total scale. The narrowness of the confidence limits associated with these coefficients indicates that they can be regarded as providing very accurate estimates of the internal consistency of the DASS in the general adult population. There is no absolute criterion for the reliability of an instrument. However, as a rule of thumb, Anastasi (1990) has suggested that  $\alpha$  should be at least .85 if the intention is to use an instrument to draw inferences concerning an individual. By this criterion all three DASS subscales and the total scale can be viewed as possessing adequate reliability.

### **Convergent and discriminant validity of the DASS**

The correlations between the anxiety and depression scales presented in Table 7, and the inferential statistical methods used to analyse them, suggest that the convergent validity of the DASS is superior to the other scales examined (e.g. the DASS scales' correlations with the other scales were significantly higher than the correlation between the other scales in three out of the four comparisons). The discriminant validity of scales is generally assessed by examining the magnitude of their correlations with measures of other constructs; a high correlation is taken as evidence of poor discriminant validity. However, in the present case there are strong theoretical grounds and empirical evidence that anxiety and depression are far from independent constructs; it is an invariant finding that such scales are highly correlated. Therefore it would have been very surprising if the DASS had bucked this trend. Moreover, the DASS scales were developed to maximize the breadth of each construct, in addition to differentiating between them. However, there was nevertheless some evidence for discriminant validity in that the within-construct correlations involving the DASS and the other self-report scales were all significantly higher than the corresponding between-construct correlations.

There is some overlap between Lovibond and Lovibond's (1995) conception of stress as measured by their stress scale, and the construct of negative affectivity. The question therefore arises whether stress is in fact equivalent to NA. The correlation between the NA and stress scales was significantly higher than NA's correlation with the other DASS scales. This is consistent with the overlap referred to above. However, the magnitude of the difference between these correlations was relatively modest (the significant effect is more a reflection of the higher statistical power conferred by the large sample size). Furthermore, it is clear from the absolute magnitude of the correlation between NA and stress (.67) that, although the constructs are associated, they cannot be viewed as interchangeable. This correlation is attenuated by measurement error in the NA and stress scales but, as both instruments are very reliable, the degree of attenuation is modest. When a correction for attenuation<sup>4</sup> was applied to the correlation (Nunnally & Bernstein, 1994), it rose to .75. Therefore, the present results indicate that, even if the constructs could be measured without error, only 56% of the variance would be shared variance.

Independent evidence that stress, as measured by the DASS, should not be regarded as synonymous with NA comes from Lovibond's (1998) study of the long-term temporal

---

<sup>4</sup>The reliability coefficients (Cronbach's alpha) used in this formula were calculated in the present sample:  $\alpha$  for the stress scale is reported in the text;  $\alpha$  for the NA scale was .85.

stability of the DASS. If the stress scale is simply an index of non-specific vulnerability to distress (i.e. NA), then stress scores at Time 1 should have been a more powerful predictor of anxiety at Time 2 than was depression, and a more powerful predictor of depression scores at Time 2 than was anxiety. Neither of these two patterns was observed, yet stress scores at Time 1 were relatively good predictors of stress scores at Time 2.

### **Conclusions and future research**

To conclude, the DASS has been shown to possess impressive psychometric properties in a large sample drawn from the general adult population. The results from CFA modelling strongly support the construct validity of the DASS scales, and the reliabilities of all three scales and the total scale were excellent. The normative data presented here should serve as useful supplements to existing normative data as they are based on a sample that was broadly representative of the general adult population in terms of age, gender and social class. The present norms are also, to our knowledge, the only UK norms currently available.

Although beyond the scope of the present investigation, it would be valuable formally to examine whether the DASS is factorially invariant. In the present study, for example, it was shown that the demographic variables (e.g. age and gender) exerted only a negligible effect on DASS scores. However, simultaneous multi-group CFA could also be employed to test whether the latent structure is invariant across age groups and gender (see Byrne, 1989, 1994). More importantly, this method could be used to examine whether the DASS is factorially invariant across cultures and across healthy and clinical populations. Examination of this latter issue would not only provide important information for those using the DASS in research or practice, but would also constitute a stringent test on the broader theoretical question of whether the constructs of anxiety, depression and stress should be viewed as continua rather than syndromes.

### **References**

- American Psychiatric Association (1994). *Diagnostic and statistical manual of mental disorders* (4th ed.). Washington, DC: American Psychiatric Association.
- Anastasi, A. (1990). *Psychological testing* (6th ed.). New York: Macmillan.
- Andrew, M. J., Baker, R. A., Kneebone, A. C., & Knight, J. L. (2000). Mood state as a predictor of neuropsychological deficits following cardiac surgery. *Journal of Psychosomatic Research*, 48, 537-546.
- Antony, M. M., Bieling, P. J., Cox, B. J., Enns, M. W., & Swinson, R. P. (1998). Psychometric properties of the 42-item and 21-item versions of the Depression Anxiety Stress Scales in clinical groups and a community sample. *Psychological Assessment*, 10, 176-181.
- Beck, A. T., Epstein, N., Brown, G., & Steer, R. A. (1988). An inventory for measuring clinical anxiety: Psychometric properties. *Journal of Consulting and Clinical Psychology*, 56, 893-897.
- Beck, A. T., Ward, C. H., Mendelsohn, M., Mock, J., & Erbaugh, J. (1961). An inventory for measuring depression. *Archives of General Psychiatry*, 4, 561-571.
- Bedford, A., & Foulds, G. (1978). *Delusions-Symptoms-States Inventory state of Anxiety and Depression*. Windsor, UK: NFER-Nelson.
- Bentler, P. (1995). *EQS structural equations program manual*. Encino, CA: Multivariate Software.

- Boyle, G. J. (1985). Self-report measures of depression: Some psychometric considerations. *British Journal of Clinical Psychology*, 24, 45-59.
- Brown, T. A., Chorpita, B. F., Korotitsch, W., & Barlow, D. H. (1997). Psychometric properties of the Depression Anxiety Stress Scales (DASS) in clinical samples. *Behaviour Research and Therapy*, 35, 79-89.
- Byrne, B. M. (1989). Multigroup comparisons and the assumption of equivalent construct validity across groups: Methodological and substantive issues. *Multivariate Behavioural Research*, 24, 503-523.
- Byrne, B. M. (1994). *Structural equation modeling with EQS and EQS/Windows*. Thousand Oaks, CA: Sage.
- Clara, I. P., Cox, B. J., & Enns, M. W. (2001). Confirmatory factor analysis of the Depression Anxiety Stress Scales in depressed and anxious patients. *Journal of Psychopathology and Behavioral Assessment*, 23, 61-67.
- Clark, L. A., & Watson, D. (1991a). Theoretical and empirical issues in differentiating depression from anxiety. In J. Becker & A. Kleinman (Eds.), *Psychosocial aspects of depression* (pp. 39-65). Hillsdale, NJ: Erlbaum.
- Clark, L. A., & Watson, D. (1991b). Tripartite model of anxiety and depression: Psychometric evidence and taxonomic implications. *Journal of Abnormal Psychology*, 100, 316-336.
- Costello, C. G., & Comrey, A. L. (1967). Scales for measuring anxiety and depression. *Journal of Psychology*, 66, 303-313.
- Feldt, L. S. (1965). The approximate sampling distribution of Kuder-Richardson Reliability Coefficient Twenty. *Psychometrika*, 30, 357-370.
- Gotlib, I. H. (1984). Depression and general psychopathology in university students. *Journal of Abnormal Psychology*, 93, 19-30.
- Horwath, E., & Weissman, M. M. (1995). Epidemiology of depression and anxiety disorders. In M. T. Tsuang, M. Tohen & G. E. P. Zahner (Eds.), *Textbook in Psychiatric Epidemiology* (pp. 317-344). New York: Wiley.
- Hu, L., & Bentler, P. M. (1999). Cutoff criteria for fit indexes in covariance structure analysis: Conventional criteria versus new alternatives. *Structural Equation Modeling*, 6, 1-55.
- Lovibond, P. F. (1998). Long-term stability of depression, anxiety, and stress syndromes. *Journal of Abnormal Psychology*, 107, 520-526.
- Lovibond, S. H., & Lovibond, P. F. (1995). *Manual for the Depression Anxiety Stress Scales*. Sydney: Psychology Foundation.
- Meltzer, H., Gill, B., Petticrew, M., & Hinds, K. (1995). *The prevalence of psychiatric morbidity among adults living in private households*. London: HMSO.
- Meng, X., Rosenthal, R., & Rubin, D. B. (1992). Comparing correlated correlation coefficients. *Psychological Bulletin*, 111, 172-175.
- Nunnally, J. C., & Bernstein, I. H. (1994). *Psychometric theory* (3rd ed.). New York: McGraw-Hill.
- Office of Population Censuses and Surveys (1990). *Classification of occupations* (1st ed., Vol. 2). London: HMSO.
- Satorra, A., & Bentler, P. M. (2001). A scaled difference chi-square test statistic for moment structure analysis. *Psychometrika*, 66, 507-514.
- Watson, D., & Clark, L. A. (1984). Negative affectivity: The disposition to experience aversive emotional states. *Psychological Bulletin*, 96, 465-490.
- Watson, D., Clark, L. A., & Tellegen, A. (1988). Development and validation of brief measures of positive and negative affect: The PANAS Scales. *Journal of Personality and Social Psychology*, 47, 1063-1070.
- Watson, D., Clark, L. A., Weber, K., Assenheimer, J. S., Strauss, M. E., & McCormick, R. A. (1995). Testing a tripartite model: II. Exploring the symptom structure of anxiety and depression in student, adult, and patient samples. *Journal of Abnormal Psychology*, 104, 15-25.
- Watson, D., & Tellegen, A. (1985). Toward a consensual structure of mood. *Psychological Bulletin*, 98, 219-235.
- Watson, D., Weber, K., Assenheimer, J. S., Clark, L. A., Strauss, M. E., & McCormick, R. A. (1995).

- Testing a tripartite model: I. Evaluating the convergent and discriminant validity of anxiety and depression symptom scales. *Journal of Abnormal Psychology*, 104, 3-14.
- Williams, E. J. (1959). The comparison of regression variables. *Journals of the Royal Statistical Society (Series B)*, 21, 396-399.
- Zigmond, A. S., & Snaith, R. P. (1983). The Hospital Anxiety and Depression Scale. *Acta Psychiatrica Scandinavica*, 67, 361-370.

*Received 25 May 2001; revised version received 18 January 2002*