



Severity of Posttraumatic Stress Symptoms—Child Age 11–17 *National Stressful Events Survey PTSD Short Scale (NSESSS)

Patient Name _____ DOB _____ Female _____ Other _____
 Provider Name _____ Date _____ Male _____

Please list the traumatic event that you experienced: _____
 Date of the traumatic event: _____

Instructions: People sometimes have problems after extremely stressful events or experiences. How much have you been bothered during the PAST SEVEN (7) DAYS by each of the following problems that occurred or became worse after an extremely stressful event/experience? **Please respond to each item by marking (✓ or x) one box per row.**

							Clinician Use
		Not at all	A little bit	Moderately	Quite a bit	Extremely	Item score
1.	Having “flashbacks,” that is, you suddenly acted or felt as if a stressful experience from the past was happening all over again (for example, you reexperienced parts of a stressful experience by seeing, hearing, smelling, or physically feeling parts of the experience)?	0	1	2	3	4	
2.	Feeling very emotionally upset when something reminded you of a stressful experience?	0	1	2	3	4	
3.	Trying to avoid thoughts, feelings, or physical sensations that reminded you of a stressful experience?	0	1	2	3	4	
4.	Thinking that a stressful event happened because you or someone else (who didn’t directly harm you) did something wrong or didn’t do everything possible to prevent it, or because of something about you?	0	1	2	3	4	
5.	Having a very negative emotional state (for example, you were experiencing lots of fear, anger, guilt, shame, or horror) after a stressful experience?	0	1	2	3	4	
6.	Losing interest in activities you used to enjoy before having a stressful experience?	0	1	2	3	4	
7.	Being “super alert,” on guard, or constantly on the lookout for danger?	0	1	2	3	4	
8.	Feeling jumpy or easily startled when you hear an unexpected noise?	0	1	2	3	4	
9.	Being extremely irritable or angry to the point where you yelled at other people, got into fights, or destroyed things?	0	1	2	3	4	
Total/Partial Raw Score:							
Prorated Total Raw Score: (if 1-2 items left unanswered)							
Average Total Score:							



Instructions to Clinicians

The National Stressful Events Survey PTSD Short Scale (NSESSS) is a 9-item measure that assesses the severity of posttraumatic stress disorder in children ages 11–17 following an extremely stressful event or experience. The measure was designed to be completed by the child upon receiving a diagnosis of posttraumatic stress disorder (or clinically significant posttraumatic stress disorder symptoms) and thereafter, prior to follow-up visits with the clinician. Each item asks the child receiving care to rate the severity of his or her posttraumatic stress disorder **during the past 7 days.**

Scoring and Interpretation

Each item on the measure is rated on a 5-point scale (0=Not at all; 1=A little bit; 2=Moderately; 3=Quite a bit, and 4=Extremely). The total score can range from 0 to 36, with higher scores indicating greater severity of posttraumatic stress disorder. The clinician is asked to review the score of each item on the measure during the clinical interview and indicate the raw score for each item in the section provided for “Clinician Use.” The raw scores on the 9 items should be summed to obtain a total raw score. In addition, the clinician is asked to calculate and use the **average total score**. The **average total score** reduces the overall score to a 5-point scale, which allows the clinician to think of the severity of the child’s posttraumatic stress disorder in terms of none (0), mild (1), moderate (2), severe (3), or extreme (4). The use of the average total score was found to be reliable, easy to use, and clinically useful to the clinicians in the DSM-5 Field Trials. The **average total score** is calculated by dividing the raw total score by number of items in the measure (i.e., 9).

Note: If 3 or more items are left unanswered, the total score on the measure should not be calculated. Therefore, the child should be encouraged to complete all of the items on the measure. If 1 or 2 items are left unanswered, you are asked to calculate a ~~prorated~~ prorated score. The prorated score is calculated by summing the scores of items that were answered to get a partial raw score. Multiply the partial raw score by the total number of items on the NSESSS—PTSD (i.e., 9) and divide the value by the number of items that were actually answered (i.e., 7 or 8). The formula to prorated the partial raw score to Total Raw Score is:

$$\frac{\text{_____ (Raw sum x 9)}}{\text{Number of items that were actually answered}}$$

If the result is a fraction, round to the nearest whole number.

Frequency of Use

To track changes in the severity of the child’s posttraumatic stress disorder over time, the measure may be completed at regular intervals as clinically indicated, depending on the stability of the child’s symptoms and treatment status. Consistently high scores on a particular domain may indicate significant and problematic areas for the child that might warrant further assessment, treatment, and follow-up. Your clinical judgment should guide your decision.



MIRACLE Counseling

NOTES:

The APA is offering a number of “emerging measures” for further research and clinical evaluation. These patient assessment measures were developed to be administered at the initial patient interview and to monitor treatment progress. They should be used in research and evaluation as potentially useful tools to enhance clinical decision-making and not as the sole basis for making a clinical diagnosis. Instructions, scoring information, and interpretation guidelines are provided; further background information can be found in DSM-5. The APA requests that clinicians and researchers provide further data on the instruments’ usefulness in characterizing patient status and improving patient care at <http://www.dsm5.org/Pages/Feedback-Form.aspx>.

Measure: Severity of Posttraumatic Stress Symptoms—Child Age 11–17
(National Stressful Events Survey PTSD Short Scale [NSESSS])

Rights granted: This measure can be reproduced without permission by researchers and by clinicians for use with their patients.

Rights holder: American Psychiatric Association

To request permission for any other use beyond what is stipulated above, contact: <http://www.appi.org/CustomerService/Pages/Permissions.aspx>

Kilpatrick DG, Resnick HS, Friedman, MJ. Copyright © 2013 American Psychiatric Association. All rights reserved.
This measure can be reproduced without permission by researchers and by clinicians for use with their patients.

8440 West national Ave. West Allis, WI 53227 | 414-405-1682 | www.miraclecounseling.org